

Neglected tropical disease targets must include morbidity

Article (Published Version)

Deribe, Kebede (2015) Neglected tropical disease targets must include morbidity. *Lancet Global Health*, 3 (10). e596. ISSN 2214-109X

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/67040/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Neglected tropical disease targets must include morbidity

The Royal Society of Tropical Medicine and Hygiene is holding a 1-day meeting in London, UK, on Sept 25, 2015, to discuss "The disease elimination agenda: the role of science, policy and advocacy".¹ Important meetings such as this are a welcome forum at which to discuss progress and challenges and to reflect on important milestones, particularly with regard to neglected tropical diseases (NTDs). Such NTD milestones include the London declaration² and the WHO roadmap.³ However, the success of these elimination initiatives is contingent on inclusive programming and on addressing the problem in its entirety.

Most of the targets for elimination focus on interrupting transmission and infection cycles, yet many of the NTDs cause severe morbidity including disabling lymphoedema, massive hydrocele, disfigurement, and blindness.⁴ Despite the huge burden of morbidities, there are no clear targets towards their elimination, and, with the exception of trachoma, there are no morbidity indicators to measure the success of elimination.⁵ A more inclusive approach to addressing morbidity in the elimination of NTDs should focus on the following points.

First, elimination targets should clearly include indicators related to morbidity. Such indicators should go beyond measuring access to care and should bind success to the extent of morbidity alleviation. WHO's trachoma elimination target of a prevalence of active trachoma of less than 5% among children aged 1–9 years and a prevalence of trachoma trichiasis of less than one case per 1000 population⁵ successfully combines both prevention of new infections and reduction of morbidity, and should be replicated across the different diseases.

Second, resources should be clearly committed to the morbidity management aspect of these NTDs. Funding such as the USAID's support of Helen Keller International's Morbidity Management and Disability Prevention for Blinding Trachoma and Lymphatic Filariasis Project is welcome.⁶ Given the scale of the problem, more resources to address the morbidity challenge are required. As resources are directed towards preventing new infection, equally resources should also be targeted to improving the quality of life of the people suffering from the consequences of the diseases.

Third, operational research into optimising the delivery of morbidity management services is also important. One of the challenges of scaling up such services is a dearth of evidence on how to integrate them into the existing health systems and ongoing NTD programmes. Operational research focusing on integration of services, surveillance, and barriers to the existing services will be important.

To achieve the challenges of elimination, morbidity management is essential, not optional. Strong advocacy and awareness-raising for donors is important, but the change should start from within by including morbidity targets in some of the NTD elimination targets.

I am supported by a Wellcome Trust Fellowship in Public Health and Tropical Medicine (grant number 099876). I declare that I have no competing interests.

Copyright © Deribe. Open Access article distributed under the terms of CC BY.

Kebede Deribe
kebededeka@yahoo.com

Brighton & Sussex Medical School, Brighton, UK;
and School of Public Health, Addis Ababa University,
Addis Ababa, Ethiopia

1 RSTMH. The disease elimination agenda: the role of science, policy and advocacy. <https://rstmh.org/events/disease-elimination-agenda-role-science-policy-and-advocacy> (accessed Aug 12, 2015).

2 Anon. Uniting to combat neglected tropical diseases: ending the neglect & reaching 2020 goals. http://unitingtocombatntds.org/sites/default/files/resource_file/london_declaration_on_ntds.pdf (accessed May 2, 2015).

3 WHO. Accelerating work to overcome the global impact of neglected tropical diseases: a roadmap for implementation. Geneva: World Health Organization, 2012. http://www.who.int/neglected_diseases/NTD_RoadMap_2012_Fullversion.pdf (accessed Aug 24, 2015).

4 Litt E, Baker MD, Molyneux D. Neglected tropical diseases and mental health: a perspective on comorbidity. *Trends Parasitol* 2012; **28**: 195–201.

5 WHO. Report of the eighth meeting of the WHO Alliance for the Global Elimination of Blinding Trachoma. Geneva: World Health Organization, 2004. http://www.who.int/blindness/publications/GET_8_Report.pdf (accessed Aug 12, 2015).

6 Helen Keller International. Helen Keller International awarded major grant by USAID to tackle neglected tropical diseases in Africa. <http://www.reuters.com/article/2015/03/24/idUSnCCN9cV9Pw+1ca+MKW20150324> (accessed Aug 12, 2015).

